

Heifermaker W/ Clarifly®

MEDICATED

Type C Medicated Feed
FOR SHOW HEIFERS AND STEERS IN CONFINEMENT
FOR CATTLE 400 LBS TO SHOW

For the prevention of coccidiosis in cattle caused by Eimeria Bovis and E. zuernii. An insect growth regulator, which prevents the development of house, stable, face and horn flies in the manure of treated beef and dairy cattle.

Active Drug Ingredients

GUARANTEED ANALYSIS

Crude Fat (min)	3.0%
Crude Fiber (max)	17%
Calcium (min)	0.5%
Calcium (max)	1.0%
Phosphorous (min)	0.20%
Salt (min)	0.20%
Salt (max)	0.40%
Potassium (min)	0.40%
Vitamin A (min)	5,000 III/lb

INGREDIENTS

Grain products, roughage products (25%), processed grain by-products, cane molasses, urea, calcium carbonate, salt, forage products, dried aspergillus oryzae fermentation product, dried aspergillus niger fermentation product, zinc amino acid complex, manganese amino acid complex, copper amino acid complex, cobalt glucoheptonate, silicon dioxide, calcium stearate, diatomaceous earth, mineral oil, zinc hydroxychloride, manganese chloride, vitamin e supplement, dextrose, lactose, colored with iron oxide, basic copper chloride, vitamin a supplement, natural flavor, artificial flavor, cobalt carbonate, ethylenediamine dihydroiodide, vitamin D3 supplement, sodium selenite.

SEE REVERSE SIDE FOR DIRECTIONS FOR USE AND PRECAUTIONARY STATEMENTS.

Guaranteed by La Grange Farm and Ranch, Inc. P.O. Box 797 • 623 E Colorado St La Grange, Texas 78945 (979) 968-6441

XXXXXX

8001



FEEDING DIRECTIONS

Feed at the rate of 2 lbs. per 100lbs. of body weight per day to provide 22.7 mg of Decoquinate per 100 pounds of body weight (0.5mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

USE DIRECTIONS FOR ACTIVE INGREDIENT DIFLUBENZURON

When used as directed, Heifermaker w/Clarifly® will provide sufficient diflubenzuron larvicide to prevent the emergence of adult house, stable, horn, and face flies from the manure of treated cattle. Existing adult flies will not be affected. Use when feed is formulated for animals 201 lbs. and over: Feeding rate will supply diflubenzuron at a level of 0.1mg per kg body weight per day. Feeding rate will supply a sufficient level of diflubenzuron to control listed flies when fed on a daily basis. Provide good quality forage and clean drinking water at all times.

CAUTION

Bentonite should not be used in decoquinate feeds. DO NOT FEED TO SHEEP DUE TO HIGH LEVELS OF SUPPLEMENTAL COPPER. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss, or death. Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

USE ONLY AS DIRECTED.

WARNING

Do not feed to cows producing milk for human consumption.



Clarifly is a product of Central Life Services.